

Minimum Cost Homomorphism Dichotomy for Oriented Cycles

Gregory Gutin¹, Arash Rafiey², Anders Yeo¹

¹ Department of Computer Science Royal Holloway, University of London, Egham, TW20 0EX, UK. e-mail: gutin@cs.rhul.ac.uk; anders@cs.rhul.ac.uk

² School of Computing Science, Simon Fraser University, Burnaby, B.C., Canada, V5A 1S6. e-mail: arashr@cs.sfu.ca

Abstract. For digraphs D and H , a mapping $f : V(D) \rightarrow V(H)$ is a homomorphism of D to H if $uv \in A(D)$ implies $f(u)f(v) \in A(H)$. If, moreover, each vertex $u \in V(D)$ is associated with costs $c_i(u)$, $i \in V(H)$, then the cost of the homomorphism f is $\sum_{u \in V(D)} c_{f(u)}(u)$. For each fixed digraph H , we have the *minimum cost homomorphism problem for H* (abbreviated $\text{MinHOM}(H)$). The problem is to decide, for an input graph D with costs $c_i(u)$, $u \in V(D)$, $i \in V(H)$, whether there exists a homomorphism of D to H and, if one exists, to find one of minimum cost. We obtain a dichotomy classification for the time complexity of $\text{MinHOM}(H)$ when H is an oriented cycle. We conjecture a dichotomy classification for all digraphs with possible loops.

Key words. Directed graph, Homomorphism, Minimum cost, Dichotomy.

1. Introduction

For directed (undirected) graphs G and H , a mapping $f : V(G) \rightarrow V(H)$ is a *homomorphism of G to H* if uv is an arc (edge) implies that $f(u)f(v)$ is an arc (edge). Let H be a fixed directed or undirected graph. The *homomorphism problem for H* asks whether a directed or undirected input graph G admits a homomorphism to H . The *list homomorphism problem for H* asks whether a directed or undirected input graph G with lists (sets) $L_u \subseteq V(H)$, $u \in V(G)$ admits a homomorphism f to H in which $f(u) \in L_u$ for each $u \in V(G)$.

Suppose G and H are directed (or undirected) graphs, and $c_i(u)$, $u \in V(G)$, $i \in V(H)$ are nonnegative costs. The cost of a homomorphism f of G to H is $\sum_{u \in V(G)} c_{f(u)}(u)$. If H is fixed, the *minimum cost homomorphism problem*, $\text{MinHOM}(H)$, for H is the following optimization problem. Given an input graph G , together with costs $c_i(u)$, $u \in V(G)$, $i \in V(H)$, we wish to find a minimum cost homomorphism of G to H , or state that none exists.

The minimum cost homomorphism problem was introduced in [10], where it was motivated by a real-world problem in defence logistics. We believe it offers a practical and natural model for optimization of weighted homomorphisms. The

problem's special cases include the homomorphism and list homomorphism problems [15, 17] and the general optimum cost chromatic partition problem, which has been intensively studied [13, 19, 20].

There is an extensive literature on the minimum cost homomorphism problem, e.g., see [5–10]. These and other papers study the time complexity of $\text{MinHOM}(H)$ for various families of directed and undirected graphs. In particular, Gutin, Hell, Rafiey and Yeo [6] proved a dichotomy classification for all undirected graphs (with possible loops): If H is a reflexive proper interval graph or a proper interval bigraph, then $\text{MinHOM}(H)$ is polynomial time solvable; otherwise, $\text{MinHOM}(H)$ is NP-hard. It is an open problem whether there is a dichotomy classification for the complexity of $\text{MinHOM}(H)$ when H is a digraph with possible loops. We conjecture that such a classification exists and, moreover, the following assertion holds:

Conjecture 1.1. Let H be a digraph with possible loops. Then $\text{MinHOM}(H)$ is polynomial time solvable if H has either a Min-Max ordering or a k -Min-Max ordering for some $k \geq 2$. Otherwise, $\text{MinHOM}(H)$ is NP-hard.

For the definitions of a Min-Max and k -Min-Max ordering see Section 3, where we give theorems (first proved in [10, 9]) showing that if H has one of the two orderings, then $\text{MinHOM}(H)$ is polynomial time solvable. So, it is the NP-hardness part of Conjecture 1.1 which is the ‘open’ part of the conjecture.

Recently Gupta, Hell, Karimi and Rafiey [5] obtained a dichotomy classification for all reflexive digraphs that confirms this conjecture. They proved that if a reflexive digraph H has no Min-Max ordering, then $\text{MinMax}(H)$ is NP-hard. Gutin, Rafiey and Yeo [8, 9] proved that if a semicomplete multipartite digraph H has neither Min-Max ordering nor k -Min-Max ordering, then $\text{MinMax}(H)$ is NP-hard.

In this paper, we show that the same result (as for semicomplete multipartite digraphs) holds for oriented cycles. This provides a further support for Conjecture 1.1. In fact, we prove a graph-theoretical dichotomy for the complexity of $\text{MinMax}(H)$ when H is an oriented cycle. The fact that Conjecture 1.1 holds for oriented cycles follows from the proof of the graph-theoretical dichotomy. In the proof, we use a new concept of a (k, l) -Min-Max ordering introduced in Section 3. Our motivation for Conjecture 1.1 partially stems from the fact that we initially proved polynomial time solvability of $\text{MinHOM}(H)$ when $V(H)$ has a (k, l) -Min-Max ordering by reducing it to the minimum cut problem. However, we later proved that (k, l) -Min-Max orderings can simply be reduced to p -Min-Max orderings for $p \geq 1$ (see Section 3).

Homomorphisms to oriented cycles have been investigated in a number of papers. Partial results for the homomorphism problem to oriented cycles were obtained in [11] and [18]. A full dichotomy was proved by Feder [3]. Feder, Hell and Rafiey [4] obtained a dichotomy for the list homomorphism problem for oriented cycles. Notice that our dichotomy is different from the ones in [3] and [4].

Bulatov [2] proved that there exists a dichotomy classification for the list homomorphism problem for digraphs, but no such dichotomy has been obtained and even conjectured to the best of our knowledge. For the homomorphism problem for

digraphs, we do not even know whether a dichotomy exists and there is no conjecture of such a classification for the general case.

The rest of this paper is organized as follows. In the next section we consider so-called levels of vertices in oriented paths and cycles. The concepts of Min-Max ordering, k -Min-Max ordering and (k, l) -Min-Max ordering are considered in Section 3. In Section 4 we obtain a dichotomy classification for $\text{MinHOM}(H)$ when H is a balanced oriented cycle. For all oriented cycles H , a dichotomy is proved in Section 5.

2. Levels of Vertices in Oriented Paths and Cycles

In this paper $[p]$ denotes the set $\{1, 2, \dots, p\}$. Let D be a digraph. We will use $V(D)$ ($A(D)$) to denote the vertex (arc) set of D . We say that xy ($x, y \in V(D)$) is an *edge* of D if either xy or yx is an arc of D . A sequence $b_1b_2 \dots b_p$ of distinct vertices of D is an *oriented path* if b_ib_{i+1} is an edge for every $i \in [p-1]$. If $b_1b_2 \dots b_p$ is an oriented path, we call $C = b_1b_2 \dots b_pb_1$ an *oriented cycle* if b_pb_1 is an edge. An edge b_ib_{i+1} (here $b_pb_{p+1} = b_pb_1$) of an oriented path P or cycle C is called *forward* (*backward*) if $b_ib_{i+1} \in A(D)$ ($b_{i+1}b_i \in A(D)$).

Let $P = b_1b_2 \dots b_p$ be an oriented path. We assign *levels* to the vertices of P as follows: we set $\text{level}_P(b_1) = 0$, and $\text{level}_P(b_{t+1}) = \text{level}_P(b_t) + 1$, if b_tb_{t+1} is forward and $\text{level}_P(b_{t+1}) = \text{level}_P(b_t) - 1$, if b_tb_{t+1} is backward. We say that P is of *type* r if $r = \max\{\text{level}_P(b_i) : i \in [p]\} = \text{level}_P(b_p)$ and $0 \leq \text{level}_P(b_t) \leq r$ for each $t \in [p]$.

An oriented cycle C is *balanced* if the number of forward edges equals the number of backward edges; if C is not balanced, it is called *unbalanced*. Note that the fact whether C is balanced or unbalanced does not depend on the choice of the vertex b_1 or the direction of C .

Let $C = b_1b_2 \dots b_pb_1$ be an oriented cycle. It has two *directions*: $b_1b_2 \dots b_pb_1$ and $b_1b_pb_{p-1} \dots b_1$. In what follows, we will always consider the direction in which the number of forward arcs is no smaller than the number of backward arcs. We can assign *levels* to the vertices of C as follows: $\text{level}(b_1) = k$, where k is a non-negative integer, and $\text{level}(b_{t+1}) = \text{level}(b_t) + 1$, if b_tb_{t+1} is forward and $\text{level}(b_{t+1}) = \text{level}(b_t) - 1$, if b_tb_{t+1} is backward. Clearly, the value of each $\text{level}(b_i)$, $i \in [p]$, depends on both k and the choice of the initial vertex b_1 . Feder [3] proved the following useful result.

Proposition 2.1. *The integer k and initial vertex b_1 in an oriented cycle C can be chosen such that $\text{level}(b_1) = 0$ and $\text{level}(b_i) \geq 0$ for every $i \in [p]$. If C is unbalanced, then k and b_1 can be chosen such that $\text{level}(b_1) = 0$ and $\text{level}(b_i) > 0$ for every $i \in [p] \setminus \{1\}$.*

Since the proposition was proved in [3], we will not give its complete proof. Instead, we will outline a procedure for finding appropriate k and b_1 and remark on how the procedure can be used in showing the proposition.

Let $C = b_1b_2 \dots b_pb_1$ be an oriented cycle. We may assume that b_1 is chosen in such a way that if C has a backward edge, then b_pb_1 is a backward edge. Compute




Fig. 1. A level diagram of oriented cycle $Z = b_1b_2 \dots b_{14}b_1$, where $b_1b_2, b_3b_2, b_3b_4, b_5b_4, b_5b_6, b_6b_7, b_8b_7b_8b_9, b_9b_{10}, b_{11}b_{10}, b_{11}b_{12}, b_{13}b_{12}, b_{14}b_{13}, b_1b_{14}$ are arcs

m_i , the number of the forward arcs minus the number of backward arcs in the oriented path $b_1b_2 \dots b_i$, for each $i \in [p]$. Set $k = |\min\{m_i : i \in [p]\}|$. Assign the level to each vertex of C using the level definition and starting from assigning level k to b_1 . By the definition of k , the level of each vertex b_j is non-negative and there are vertices b_i of level zero. Choose such a vertex b_i with maximum index i and reassign the levels to the vertices of C as follows. Consider $C' = b_i b_{i+1} \dots b_p b_1 b_2 \dots b_i$ and set $\text{level}(b_i) = 0$ and the rest of the levels according to the order of vertices given in C' .

This procedure can be turned into a proof of the proposition by observing that if C' is unbalanced, then the level of b_1 in C' will be greater than the level of b_1 in C . Thus, the levels of all vertices v_j , $j \in [i-1]$ will be greater than their levels in C , implying that the only level zero vertex in C' is b_i .

Thus, in the rest of the paper, we may assume that the ‘first’ vertex of b_1 of an oriented cycle $C = b_1b_2 \dots b_pb_1$ is chosen in such a way that the levels of all vertices of C satisfy Proposition 2.1.

We will extensively use the following notation: $VL(C) = \{b_t : \text{level}(b_t) = 0, t \in [p]\}$, $h(C) = \max\{\text{level}(b_j) : j \in [p]\}$, and $VH(C) = \{b_t : \text{level}(b_t) = h(C), t \in [p]\}$. Note that for unbalanced cycles C we have $|VL(C)| = 1$.

The concepts of this section are illustrated on Figure 1. In particular, Z is balanced with forward edges b_1b_2, b_3b_4, \dots and backward edges b_2b_3, b_4b_5, \dots . We have $\text{level}(b_1) = \text{level}(b_3) = \text{level}(b_5) = 0$, $\text{level}(b_2) = \text{level}(b_4) = \text{level}(b_6) = \text{level}(b_8) = \text{level}(b_{14}) = 1$, $\text{level}(b_7) = \text{level}(b_9) = \text{level}(b_{11}) = \text{level}(b_{13}) = 2$ and $\text{level}(b_{10}) = \text{level}(b_{12}) = 3$. Thus, $h(Z) = 3$, $VL(Z) = \{b_1, b_3, b_5\}$, $VH(Z) = \{b_{10}, b_{12}\}$.

3. k -Min-Max and (k, l) -Min-Max Orderings

All known polynomial cases of $\text{MinHOM}(H)$ can be formulated in terms of certain vertex orderings. In fact, all known polynomial cases can be partitioned into two

classes: digraphs H admitting a Min-Max ordering of their vertices and digraphs H having a k -Min-Max ordering of their vertices ($k \geq 2$). Both types of orderings are defined in this section, where we also introduce a new type of ordering, a (k, l) -Min-Max ordering. It may be surprising, but we prove that the new type of ordering can be reduced to the two known orderings.

Let H be a digraph and let (v_1, v_2, \dots, v_p) be an ordering of the vertices of H . Let $e = v_i v_r$ and $f = v_j v_s$ be two arcs in H . The pair $v_{\min\{i,j\}} v_{\min\{s,r\}} (v_{\max\{i,j\}} v_{\max\{s,r\}})$ is called the *minimum (maximum)* of the pair e, f . (The minimum (maximum) of two arcs is not necessarily an arc.) An ordering (v_1, v_2, \dots, v_p) is a *Min-Max ordering* of $V(H)$ if both minimum and maximum of every two arcs in H are in $A(H)$. Two arcs $e, f \in A(H)$ are called a *crossing pair* if $\{e, f\} \neq \{g', g''\}$, where g' (g'') is the minimum (maximum) of e, f . Clearly, to check that an ordering is Min-Max, it suffices to verify that the minimum and maximum of every crossing pair of arcs are arcs, too. The concept of Min-Max ordering is of interest due to the following:

Theorem 3.1. [10] *If a digraph H has a Min-Max ordering of $V(H)$, then $\text{Min-HOM}(H)$ is polynomial-time solvable.*

We will sometimes call a Min-Max ordering also a *1-Min-Max ordering*. The reason for this will become apparent in the rest of this section.

A collection V_1, V_2, \dots, V_k of subsets of a set V is called a *k-partition* of V if $V = V_1 \cup V_2 \cup \dots \cup V_k$, $V_i \cap V_j = \emptyset$ provided $i \neq j$.

Let $H = (V, A)$ be a digraph and let $k \geq 2$ be an integer. We say that H has a *k-Min-Max ordering* of $V(H)$ if there is a k -partition of V into subsets V_1, V_2, \dots, V_k and there is an ordering $\vec{V}_i = (v_1^i, v_2^i, \dots, v_{\ell(i)}^i)$ of V_i for each i such that

- (i) Every arc of H is an arc from V_i to V_{i+1} for some $i \in [k]$ and
- (ii) $(\vec{V}_i, \vec{V}_{i+1}) = (v_1^i, v_2^i, \dots, v_{\ell(i)}^i v_1^{i+1} v_2^{i+1}, \dots, v_{\ell(i+1)}^{i+1})$ is a Min-Max ordering of the subdigraph of H induced by $V_i \cup V_{i+1}$ for each $i \in [k]$. (All indices are taken modulo k .)

In such a case, $(\vec{V}_1, \vec{V}_2, \dots, \vec{V}_k)$ is a *k-Min-Max ordering* of $V(H)$; k -Min-Max orderings are of interest due to the following:

Theorem 3.2. [9] *If a digraph H has a k-Min-Max ordering of $V(H)$, then $\text{Min-HOM}(H)$ is polynomial-time solvable.*

Our study of $\text{MinHOM}(H)$ for oriented cycles H has led us to the following new concept.

Definition 3.3. Let $H = (V, A)$ be a digraph and let $k \geq 2$ and l be integers. For $l < k$ we say that H has a *(k, l)-Min-Max ordering* if there is a $(k + l - 2)$ -partition of V into subsets $V_1, V_2, \dots, V_k, U_2, U_3, \dots, U_{l-1}$ (set $U_1 = V_1, U_l = V_k$) and there is an ordering $\vec{V}_i = (v_1^i, v_2^i, \dots, v_{\ell^v(i)}^i)$ of V_i for each $1 \leq i \leq k$ and there is an ordering $\vec{U}_i = (u_1^i, u_2^i, \dots, u_{\ell^u(i)}^i)$ of U_i for each $1 \leq i \leq l$ such that

- (i) Every arc of H is an arc from V_i to V_{i+1} for some $i \in [k-1]$, or is an arc from U_j to U_{j+1} for some $j \in [l-1]$.
- (ii) $(\vec{V}_i, \vec{V}_{i+1})$ is a Min-Max ordering of the subdigraph $H[V_i \cup V_{i+1}]$ for all $i \in [k-1]$.
- (iii) $(\vec{U}_i, \vec{U}_{i+1})$ is a Min-Max ordering of the subdigraph $H[U_i \cup U_{i+1}]$ for all $i \in [l-1]$.
- (iv) $(\vec{V}_1, \vec{V}_2, \vec{U}_2)$ is a Min-Max ordering of the subdigraph $H[V_1 \cup V_2 \cup U_2]$.
- (v) $(\vec{U}_{l-1}, \vec{V}_{k-1}, \vec{V}_k)$ is a Min-Max ordering of the subdigraph $H[V_{k-1} \cup U_{l-1} \cup V_k]$.

It turns out that (k, l) -Min-Max orderings can be reduced to p -Min-Max orderings as follows from the next assertion:

Theorem 3.4. *If a digraph H has a (k, l) -Min-Max ordering, then $\text{MinHOM}(H)$ is polynomial-time solvable.*

Proof. Let H have a (k, l) -Min-Max ordering as described in Definition 3.3. Let $d = k - l$. We will show that H has a d -Min-Max ordering, which will be sufficient because of Theorems 3.1 and 3.2. (Recall that a 1-Min-Max ordering is simply a Min-Max ordering.) Let us consider two cases.

Case 1. $d = 1$. It is not difficult to show that the ordering

$$(\vec{V}_1, \vec{V}_2, \vec{U}_2, \vec{V}_3, \vec{U}_3, \dots, \vec{V}_{k-2}, \vec{U}_{k-2}, \vec{V}_{k-1}, \vec{V}_k)$$

is a Min-Max ordering. Indeed, all crossing pairs of arcs are only in the subgraphs given in (ii)-(v) of Definition 3.3. According to the definition, the maximum and minimum of every crossing pair is in H .

Case 2. $d \geq 2$. Let $s_i = \max\{p : i + pd \leq l\}$ for each $i \in [d]$. Consider the following orderings for each $2 \leq i \leq d$

$$\vec{W}_i = (\vec{V}_i, \vec{U}_i, \vec{V}_{i+d}, \vec{U}_{i+d}, \vec{V}_{i+2d}, \vec{U}_{i+2d}, \dots, \vec{V}_{i+s_id}, \vec{U}_{i+s_id}, \vec{V}_{i+(s_i+1)d})$$

and the ordering

$$\vec{W}_1 = (\vec{V}_1, \vec{V}_{1+d}, \vec{U}_{1+d}, \vec{V}_{1+2d}, \vec{U}_{1+2d}, \dots, \vec{V}_{1+s_1d}, \vec{U}_{1+s_1d}, \vec{V}_{1+(s_1+1)d}).$$

Observe that W_1, W_2, \dots, W_d form a partition of $V(H)$ and that every arc is from W_i to W_{i+1} for some $i \in [d]$, where $W_{d+1} = W_1$. As in Case 1, it is not difficult to see that $(\vec{W}_1, \vec{W}_2, \dots, \vec{W}_d)$ is a d -Min-Max ordering of H . \square

Remark 3.5. Notice that not always a p -Min-Max ordering can be reduced to a (k, l) -Min-Max ordering. As an example, consider \vec{C}_5 .

4. Balanced Oriented Cycles

We say that a balanced oriented cycle $C = b_1 b_2 \dots b_p b_1$ is of the form $(l^+ h^+)^q$ with $q \geq 1$ if $P = C - b_p b_1$ can be written as $P = x_1 P_1 y_1 R_1 x_2 P_2 y_2 R_2 \dots x_q P_q y_q R_q$, where $x_i \in VL(C)$, $y_i \in VH(C)$ for each $i \in [q]$, P_i, R_i are oriented paths and all vertices in $VL(C) \cap V(P_i)$ ($VH(C) \cap V(R_i)$) appear before all vertices in $VH(C) \cap V(P_i)$ in P_i ($VL(C) \cap V(R_i)$ in R_i) for each $i \in [q]$. We write $l^+ h^+$ instead of $(l^+ h^+)^1$. For example, the cycle in Figure 1 is of the form $l^+ h^+$. Balanced oriented cycles C of the form $l^+ h^+$ are considered in the following:

Theorem 4.1. *Let $C = b_1 b_2 \dots b_p b_1$ be a balanced oriented cycle of the form $l^+ h^+$. Then $\text{MinHOM}(C)$ is polynomial time solvable.*

Proof. Let $q = \min\{j : b_j \in VH(C)\}$, let $m = h(C)$ and let $V = V(C)$. Consider the following ordering $\vec{V} = (b_p, b_{p-1}, \dots, b_{q+1}, b_1, b_2, \dots, b_q)$ of V . We can define the following natural $(m+1)$ -partition of V : V_1, V_2, \dots, V_{m+1} , where $V_j = \{b_s \in V(C) : \text{level}(b_s) = j-1\}$. Note that every arc of C is an arc from V_j to V_{j+1} for some $j \in [m]$.

Let \vec{V}_j be the ordering of V_j obtained from \vec{V} by deleting all vertices not in V_j and let $\vec{V}_j = (s_1^j, s_2^j, \dots, s_{b(j)}^j)$. Observe that the digraph $C[V_j \cup V_{j+1}]$ has no crossing pair of arcs for any $j \in [m]$ since, for every pair $s_\alpha^j s_\beta^{j+1}, s_\gamma^j s_\delta^{j+1}$ of arcs in the digraph, we have that either $\alpha \leq \gamma$ and $\beta \leq \delta$, or $\alpha \geq \gamma$ and $\beta \geq \delta$. Thus, C has an $(m+1)$ -Min-Max ordering of vertices and, by Theorem 3.2, $\text{MinHOM}(C)$ is polynomial-time solvable. \square

The following lemma was first proved in [12]; see also [3, 21] and Lemma 2.36 in [17].

Lemma 4.2. *Let P_1 and P_2 be two oriented paths of type r . Then there is an oriented path P of type r that maps homomorphically to P_1 and P_2 such that the initial vertex of P maps to the initial vertices of P_1 and P_2 and the terminal vertex of P maps to the terminal vertices of P_1 and P_2 . The length of P is polynomial in the lengths of P_1 and P_2 .*

We need a modified version of Lemma 4.2, Lemma 4.3. We say that an oriented path $b_1 b_2 \dots b_p$ of type r is of the form $(l^+ h^+)^k$ if the balanced oriented cycle

$$b_1 b_2 \dots b_p a_{r-1} a_{r-2} \dots a_2 a_1 b_1$$

is of the form $(l^+ h^+)^k$, where $b_1 a_1 a_2 \dots a_{r-2} a_{r-1} b_p$ is a directed path.

Lemma 4.3. *Let P_1 and P_2 be two oriented paths of type r . Let P_1 be of the form $h^+ l^+$ and let P_2 be of the form $(l^+ h^+)^k$, $k \geq 1$. Then there is an oriented path P of type r that maps homomorphically to P_1 and P_2 such that the initial vertex of P maps to the initial vertices of P_1 and P_2 and the terminal vertex of P maps to the terminal vertices*

of P_1 and P_2 . The length of P is polynomial in the lengths of P_1 and P_2 , and P is of the form $(l^+h^+)^k$.

Proof. We will show that our construction implies that $|V(P)| \leq |V(P_1)| \times |V(P_2)|$.

We first prove the lemma for the case when $k = 1$. The proof is by induction on $r \geq 0$. If $0 \leq r \leq 1$, the claim is trivial. Assume that $r \geq 2$. Let $P_1 = a_1a_2 \dots a_p$, let $P_2 = b_1b_2 \dots b_q$, let $s_1 = \min\{i : \text{level}_{P_1}(a_i) = r\}$ and let $s_2 = \min\{i : \text{level}_{P_2}(b_i) = r\}$. Let $\beta_1 = \min\{\text{level}_{P_1}(a_i) : s_1 \leq i\}$ and $\beta_2 = \min\{\text{level}_{P_2}(b_i) : s_2 \leq i\}$. Without loss of generality assume that $\beta_1 \leq \beta_2$ and let $t_1 = \min\{i : \text{level}_{P_1}(a_i) = \beta_1 \text{ and } i \geq s_1\}$ and let $s_2 = \max\{i : \text{level}_{P_2}(b_i) = \beta_1 \text{ and } i \leq s_2\}$. Note that $\beta_1 > 1$ as P_1 is of form h^+l^+ .

By the induction hypothesis, there is an appropriate oriented path P' that can be mapped homomorphically to $a_1a_2 \dots a_{s_1-1}$ and $b_1b_2 \dots b_{s_2-1}$. There is also an oriented path P'' that can be mapped homomorphically to $a_{s_1}a_{s_1+1} \dots a_{t_1}$ and $b_{s_2}b_{s_2+1} \dots b_{t_2}$ (by reversing the two paths and then reversing the path we get by the induction hypothesis). Furthermore there is an oriented path P''' that can be mapped homomorphically to $a_{t_1+1}a_{t_1+2} \dots a_p$ and $b_{t_2+1}b_{t_2+2} \dots b_q$. Let $P = P'P''P'''$ (where the arc between the last vertex of P' to the first vertex of P'' is oriented from P' to P'' and similarly the arc between P'' and P''' is oriented in that direction). Note that P is of type r and form h^+l^+ and maps homomorphically to P_1 and P_2 such that the initial vertex of P maps to the initial vertices of P_1 and P_2 and the terminal vertex of P maps to the terminal vertices of P_1 and P_2 . Furthermore $|V(P)| \leq (s_1 - 1)(s_2 - 1) + (t_1 - s_1 + 1)(s_2 - t_2 + 1) + (p - s_1)(q - t_2)$. As $(s_1 - 1) + (t_1 - s_1 + 1) + (p - s_1) = p$ and $(s_2 - 1) + (s_2 - t_2 + 1) + (q - t_2) \leq q$ we have $|V(P)| \leq pq \leq |V(P_1)| \times |V(P_2)|$.

Now we proceed by induction on $k \geq 1$. The base case has already been proved. Assume that $k \geq 2$ and let $P_1 = a_1a_2 \dots a_p$ and $P_2 = b_1b_2 \dots b_q$. Let $t = \max\{i : \text{level}_{P_2}(b_i) = 0\}$ and let $s = \max\{i : i < t, \text{level}_{P_2}(b_i) = r\}$. By the induction hypothesis, there is an appropriate oriented path P' that can be mapped homomorphically to P_1 and $b_1b_2 \dots b_s$. Also, there is an appropriate oriented path P'' (P''') that can be mapped homomorphically to $a_p a_{p-1} \dots a_1$ and $b_s b_{s+1} \dots b_t$ (P_1 and $b_t b_{t+1} \dots b_q$). Now obtain a new oriented path P by identifying the terminal vertex of P' with the initial vertex of P'' and the terminal vertex of P'' with the initial vertex of P''' . Observe that P satisfies the required properties. \square

Consider the oriented cycle $C_4^0 = 12341$ with arcs 12, 32, 14, 34. Observe that 1, 2, 3, 4 is a Min-Max ordering of $V(C_4^0)$ and, thus, $\text{MinHOM}(C_4^0)$ is polynomial-time solvable.

Theorem 4.4. Let $C = b_1b_2 \dots b_p b_1$ be a balanced oriented cycle of the form $(l^+h^+)^k$, $k \geq 2$, and let $C \neq C_4^0$. Then $\text{MinHOM}(C)$ is NP-hard.

Proof. Let $C \neq C_4^0$. Let $s = \min\{j : b_j \in VH(C)\}$, $q = \min\{j : j > s, b_j \in VL(C)\}$, $t = \min\{j : j > q, b_j \in VH(C)\}$ and $m = h(C)$. Let $P_1 = b_1b_2 \dots b_s$, $P_2 = b_q b_{q-1} \dots b_s$, $P_3 = b_q b_{q+1} \dots b_t$ and $P_4 = b_1 b_p b_{p-1} \dots b_t$. Note that each

P_j is of type m . By Lemma 4.3 there is a path Q_1 of type m which is mapped homomorphically to P_4 , P_2 and P_3 . There is also a path Q_2 of type m and which is mapped homomorphically to P_1 and P_3 . Since $C \neq C_4^0$ and the end-vertices of Q_1 are mapped to the end-vertices of P_4 , the path Q_1 contains more than two vertices. Furthermore, by Lemma 4.3 we may assume that Q_1 is of the form $(l^+h^+)^{k-1}$ and Q_2 is of the form l^+h^+ .

Let x (y) be the terminal vertex of Q_1 (Q_2). Form a new oriented path $Q = q_1q_2 \dots q_l$ by identifying x with y and let $1 \leq r \leq l$ be defined such that $Q_1 = q_1q_2 \dots q_r$ and $Q_2 = q_lq_{l-1} \dots q_r$. As Q_1 contains more than two vertices we have that $r \geq 3$.

Let D be an arbitrary digraph. We will now reduce the problem of finding a maximum independent set in D (i.e. in the underlying graph of D) to $\text{MinHOM}(C)$. Replace every arc ab of D by a copy of Q identifying q_1 with a and q_l with b , and denote the obtained digraph by D' . For every path Q in D' which we added in the construction of D' we define the cost function c as follows, where M is a number greater than $|V(D)|$:

- (i) $c_{b_1}(q_1) = c_{b_1}(q_l) = 0$ and $c_{b_q}(q_1) = c_{b_q}(q_l) = 1$;
- (ii) $c_{b_s}(q_r) = c_{b_t}(q_r) = 0$ and $c_b(q_r) = M$ for all $b \in V(C) - \{b_s, b_t\}$;
- (iii) $c_{b_2}(q) = M$ for all $q \in \{q_2, q_3, \dots, q_{r-1}\} (\neq \emptyset, \text{ as } r \geq 3)$;
- (iv) if $s = 2$ then $c_{b_1}(q_{r-1}) = M$;
- (v) all other costs of mapping vertices of D' to H are zero.

Consider a mapping g from $V(Q)$ to $V(C)$, where $g(q_1) = b_q$, $g(q_l) = b_q$, and Q_1 and Q_2 are both homomorphically mapped to P_3 (i.e., $g(q_r) = b_t$). Observe that g is a homomorphism from Q to C of cost 2. This implies, in particular, that there is a homomorphism from D' to H of cost less than M . We now consider three other homomorphisms from Q to C :

- (a) $f(q_1) = b_1$ and $f(q_l) = b_q$, and Q_1 is mapped to P_4 and Q_2 is mapped to P_3 homomorphically (i.e., $f(q_r) = b_t$). The cost of f is 1.
- (b) $f'(q_1) = b_q$ and $f'(q_l) = b_1$, and f' maps Q_1 to P_2 and Q_2 to P_1 homomorphically (i.e., $f'(q_r) = b_s$). The cost of f' is 1.
- (c) $f''(q_1) = b_1$ and $f''(q_l) = b_1$. We will show that the cost of f'' is at least M . If this is not the case then $f''(q_r) \in \{b_s, b_t\}$ by (ii). First assume that $f''(q_r) = b_s$. By (iii) no vertex of $V(Q_1) - \{q_r\}$ is mapped to b_2 and if $s = 2$ then by (iv) q_{r-1} is not mapped to b_1 . However as Q_1 is of the form $(l^+h^+)^{k-1}$ and the path $b_1b_p b_{p-1} \dots b_s$ is of the form $(l^+h^+)^k$ we get a contradiction to f'' mapping Q_1 to C and $f''(q_1) = b_1$ and $f''(q_r) = b_s$. So now assume that $f''(q_r) = b_t$. However Q_2 is of the form (l^+h^+) and there is no path in C from b_1 to b_t of the form (l^+h^+) , a contradiction. Therefore the cost of f'' is at least M .

By the above a minimum cost homomorphism $h : D' \rightarrow C$ maps all vertices from a maximum independent set in D to b_1 and all other vertices from D to b_q . As finding a maximum independent set in a digraph is NP-hard we see that $\text{MinHOM}(C)$ is NP-hard. \square

5. Dichotomy and Unbalanced Oriented Cycles

We are ready to prove the following main result:

Theorem 5.1. *Let C be an oriented cycle. If C is unbalanced or C is balanced of the form l^+h^+ or $C = C_4^0$, then $\text{MinHOM}(C)$ is polynomial-time solvable. Otherwise, $\text{MinHOM}(C)$ is NP-hard.*

By Theorems 4.1 and 4.4, to show Theorem 5.1 it suffices to prove the following:

Theorem 5.2. *Let $C = b_1b_2 \dots b_pb_1$ be an unbalanced oriented cycle. Then $\text{MinHOM}(C)$ is polynomial-time solvable.*

Proof. It is well-known that the minimum cost homomorphism problem to a directed cycle is polynomial-time solvable (see, e.g., [7]). Thus, we may assume that C is not a directed cycle. By Proposition 2.1, we may assume that $\text{level}(b_1) = 0$ and $\text{level}(b_i) > 0$ for all $i \in [p] \setminus \{1\}$. Let $q = \max\{j : b_j \in V H(C)\}$.

Consider the oriented path $P = b_1b_2 \dots b_q$. Let $V_{i+1} = \{b_j \in V(P) : \text{level}(b_j) = i\}$ for all $i \in [k] \cup \{0\}$, where $k = \text{level}(b_q)$. Now consider the oriented path $Q = b_1b_pb_{p-1} \dots b_q$. Assign levels to the vertices of Q stating from $\text{level}_Q(b_1) = 0$ and continuing as described in Section 1. Observe that all vertices of Q get non-negative levels. Let $U_{i+1} = \{b_j \in V(Q) : \text{level}_Q(b_j) = i\}$, $i \in [l] \cup \{0\}$, where $l = \text{level}_Q(b_q)$. Clearly, $V_1 = U_1 = \{b_1\}$; set $U_{l+1} = V_{k+1}$.

Consider the ordering $\vec{U} = (b_1, b_p, b_{p-1}, \dots, b_q)$ of the vertices of Q . For $i \in [l+1]$, the ordering \vec{U}_i is obtained from \vec{U} by deleting all vertices not in U_i . Consider the ordering $\vec{V} = (b_1, b_2, \dots, b_q)$ of the vertices of P . For $i \in [k+1]$, the ordering \vec{V}_i is obtained from \vec{V} by deleting all vertices not in V_i . Observe that the ordering $(\vec{V}_1, \vec{V}_2, \vec{U}_2)$ of the vertices of $C[V_1 \cup V_2 \cup U_2]$ has no crossing arcs. Similarly, the ordering $(\vec{U}_l, \vec{V}_k, \vec{V}_{k+1})$ of the vertices of $C[U_l \cup V_k \cup V_{k+1}]$ has no crossing arcs, and orderings $(\vec{V}_i, \vec{V}_{i+1})$ and $(\vec{U}_j, \vec{U}_{j+1})$ ($i \in [k]$, $j \in [l]$) have no crossing arcs. Thus, C has a $(k+1, l+1)$ -ordering of vertices. Now we are done by Theorem 3.4. \square

References

1. Bang-Jensen, J., Gutin, G.: Digraphs: Theory, Algorithms and Applications. Springer-Verlag, London, (2000)
2. Bulatov, A.A.: Tractable conservative constraint satisfaction problems. In: Proceedings of 18th Annual IEEE Symposium on Logic in Computer Science, pp. 321–330, Ottawa, Canada, June (2003). IEEE Computer Society
3. Feder, T.: Homomorphisms to oriented cycles and k -partite satisfiability. SIAM J. Discrete Math **14**, 471–480 (2001)
4. Feder, T., Hell, P., Rafiey, A.: List homomorphism to balanced digraphs, Submitted for Publication
5. Gupta, A., Hell, P., Karimi, M., Rafiey, A.: Minimum cost homomorphisms to reflexive digraphs. In: Proceedings of 8th Latin American Symposium on Theoretical Informatics (LATIN'08), Lecture Notes Comput. Sci. **4957**, 182–193 (2008)

6. Gutin, G., Hell, P., Rafiey, A., Yeo, A.: Minimum cost homomorphisms to proper interval graphs and bigraphs. *Europ. J. Combin.* **29**, 900–911 (2008)
7. Gutin, G., Rafiey, A., Yeo, A.: Minimum cost and list homomorphisms to semicomplete digraphs. *Discrete Appl. Math.* **154**, 890–897 (2006)
8. Gutin, G., Rafiey, A., Yeo, A.: Minimum cost homomorphisms to semicomplete multipartite digraphs. *Discrete Appl. Math.* **156**, 2429–2435 (2008)
9. Gutin, G., Rafiey, A., Yeo, A.: Minimum cost homomorphisms to semicomplete bipartite digraphs. *SIAM J. Discrete Math.* **22**, 1624–1639 (2008)
10. Gutin, G., Rafiey, A., Yeo, A., Tso, M.: Level of repair analysis and minimum cost homomorphisms of graphs. *Discrete Appl. Math.* **154**, 881–889 (2006)
11. Gutjahr, W.: Graph Colourings, Ph.D. thesis (Free University, Berlin, 1991)
12. Häggkvist, R., Hell, P., Miller, D.J., Neumann-Lara, V.: On multiplicative graphs and the product conjecture. *Combinatorica* **8**, 63–74 (1988)
13. Halldorsson, M.M., Kortsarz, G., Shachnai, H.: Minimizing average completion of dedicated tasks and interval graphs. *Approximation, Randomization, and Combinatorial Optimization* (Berkeley, Calif, 2001), *Lecture Notes in Computer Science*, vol. **2129**, pp. 114–126 Springer, Berlin (2001)
14. Feder, T., Hell, P., Huang, J.: List homomorphisms and circular arc graphs. *Combinatorica* **19**, 487–505 (1999)
15. Hell, P.: Algorithmic aspects of graph homomorphisms. In: *Survey in Combinatorics 2003*. London Math. Soc. Lecture Note Series, vol. **307** pp. 239–276. Cambridge University Press, Cambridge (2003)
16. Hell, P., Nešetřil, J.: On the complexity of H -colouring. *J. Combin. Theory B* **48**, 92–110 (1990)
17. Hell, P., Nešetřil, J.: *Graphs and Homomorphisms*. Oxford University Press, Oxford (2004)
18. Hell, P., Zhu, X.: The existence of homomorphisms to oriented cycles. *SIAM J. Discrete Math.* **8**, 208–222 (1995)
19. Jansen, K.: Approximation results for the optimum cost chromatic partition problem. *J. Algorithms* **34**, 54–89 (2000)
20. Jiang, T., West, D.B.: Coloring of trees with minimum sum of colors. *J. Graph Theory* **32**, 354–358 (1999)
21. Zhu, X.: A simple proof of the multiplicativity of directed cycles of prime power length. *Discrete Appl. Math.* **36**, 333–345 (1992)

Received: December 22, 2007

Final Version received: February 27, 2009